

SEMPER TRAVEL 1

Class 2 OTC drug
Motion Sickness Products

◆SEMPER TRAVEL 1 prevents and alleviates symptoms of motion sickness such as dizziness, nausea, and headache. ◆Once daily oral dose takes effect, and helps your fun trip and helps you to travel comfortably. ◆The grapefruit-flavor chewable tablets can dissolve in the mouth without water, or be chewed down. ◆The drug relieves symptoms when you take it even after you begin to feel sick.

Indication

1) prevention and relief of dizziness, nausea, and headache associated with motion sickness

Dosage and administration

Take the following dose by chewing or dissolving the drug in the mouth.

To prevent motion sickness, take the drug 30 minutes prior to boarding.

15 years or over: 1 tablet per dose, once daily

7 to 14 years: 1/2 tablet per dose, once daily

Under 7 years: Do not take

Comply with the prescribed dosage and administration instructions.

Use for children should be supervised by a parent.

•As shown in the figure, press the convex part of the PTP sheet containing the tablet firmly with your fingertips, break the aluminum foil on the back side, and take out the tablets.

How to take out the tablets

• (If the tablets are inadvertently taken together with the PTP sheet containing them, it will lead to unexpected accidents, such as the sharp edge sticking into the esophageal mucosa.)

ingredient and amount

In 1 tablets

Chlorpheniramine Maleate 4mg

Scopolamine Hydrobromide 0.25mg

Excipients

Reduced maltose syrup, Corn starch, Hydroxypropylcellulose, anhydrous silicic acid, Aspartame (L- phenylalanine compound), Magnesium stearate, flavor, Starch sodium octenyl succinate

Precautions

When not to use the product

(If you do not follow these instructions, the current symptoms may worsen or adverse reactions/incidents are more likely to occur.)

This drug should not be taken together with the following drugs:

Other medicine for motion sickness, cold medicine, antipyretics/analgesics, sedatives, expectorant, indigestion and heartburn relief, internal medicine containing antihistamines, etc. (such as internal medicine for rhinitis and medicine for allergies)

●After taking this drug, do not drive a car or operate machinery (symptoms such as sleepiness, blurred vision and abnormal brightness may occur).

Consultation

●The following persons should contact a physician, pharmacist, or registered salesperson for a consultation before administration.

Patients undergoing medical treatment from a physician

Pregnant women or women suspected of being pregnant.

The elderly

Patients who have experienced allergic symptoms associated with drugs, etc.

Persons with the following symptoms:

urination difficulty

Persons diagnosed as having the following

Glaucoma

heart disease

●If the following symptoms are observed after taking this drug, these may be adverse reactions, so immediately discontinue the use of this drug, and show this document to your physician, pharmacist, or registered salesperson for a consultation.

Skin

rash/redness

itching

Neuropsychiatric system

headache

Urinary system:

urination difficulty

Other:

Facial hot flashes

abnormal brightness

The following serious symptoms may occur in rare cases. In such cases, immediately seek medical aid from a physician.

Aplastic anemia: Symptoms, such as bruises, nose bleeds, bleeding gums, fever, pale appearance of skin and mucosa, fatigue, palpitations, shortness of breath, becoming dizzy from feeling sick and blood in the urine, etc. may occur.

Agranulocytosis: Symptoms, such as acute high fever, chills and sore throat, etc. may occur.

●The following symptoms may be observed after taking this drug. If these symptoms persist or worsen, discontinue the use of this drug, and show this document to your physician, pharmacist, or registered salesperson for a consultation.

Thirst, sleepiness, constipation, blurred vision

Precautions for storage and handling

Avoid direct sunlight, and store the product in a cool place with little humidity.

Store the product beyond the reach of children.

Do not transfer the drug to other containers. (It may lead to misuse or quality deterioration)

Do not take the product past the expiration date.

12

【Disclaimer on Multilingual OTC Product Information】

•This product is a pharmaceutical product approved under a Japanese law, the Law for Ensuring the Quality, Efficacy and Safety of Drugs and Medical Devices, with a view to its sale and use in Japan.

•Multilingual product information is a translation of the product labeling written in Japanese and provided for your information only. It does not warrant that its contents and the product itself conforms to laws and regulations in countries other than Japan.

•Multilingual product information is a tentative translation by the provider (or Our Company), and may be modified or altered without notice.

•The provider (or Our Company) assumes no responsibility for any occurred problem attributable to the contents of the multilingual product information.